



West of Scotland Guideline

Approved February 2019

What's New:

Mattresses should be vacuumed
Updated BNF information

Phthirus pubis – pubic lice

The crab louse (phthirus pubis) is transmitted by close body contact. Affects coarse hair of the pubic area, body and rarely the eyebrows and eyelashes. Incubation period is usually between 5 days and several weeks.

Signs and Symptoms

- Itch
- Identifying the eggs or crab louse (tan/grey in colour) or faecal specks (black)
- Occasionally some individuals can have prolonged, asymptomatic infestation

Diagnosis

- Finding the adult lice and/or eggs seen in affected areas - a magnifying glass may help. Eggs adhere to the hair
- Blue macules (maculae caeruleae) may be visible at feeding sites
- Examination under light microscopy can confirm the exact morphology
- Nb: May also affect eyelashes and eyebrows

Management

General advice

- Avoid close body contact until they and their partner(s) have completed treatment
- Give detailed explanation of the condition, and clear and accurate written information on applying the treatment. All surfaces of the body should be treated, including the scalp, neck, and face (paying particular attention to the eyebrows and other facial hair)
- Offer STI testing

First Line Treatment:

- **Malathion 0.5% Aqueous Lotion (Derbac M)-** apply over whole body (dry) and wash off 12 hours later. Give 100ml
- Repeat after 7 days
- **Eyelashes:** Simple eye ointment BP can be applied to eyelash and eyelash root bd for 8-10 days, **this avoids any eye irritation that may occur if using an insecticide**

WOS PUBIC LICE PROTOCOL	APPROVED
WOS MCN CLINICAL GUIDELINES GROUP	VERSION: DRAFT 6.1 LAST UPDATED: Feb 19
REVIEW DATE: Feb 2021 PAGE NUMBER: 1 of 2	COPIES AVAILABLE: www.wossexualhealthmcn.org



West of Scotland Guideline

Approved February 2019

- Consider advising shaving particularly hairy areas prior to treatment
- Lotions should be applied to all body hair, including the beard and moustache and eyebrows but excluding scalp (pubic lice do not attach to soft head hair).
- Following treatment if nits are still attached to the hair shaft then these can be removed by a fingernail or a fine tooth comb
- To kill any lice or nits remaining on clothing, towels, or bedding, machine-wash and machine-dry those items that the infested person used during the 2-3 days before treatment. Use hot water (at least 50°C) and the hot dryer cycle.
- Items that cannot be laundered can be dry-cleaned or stored in a sealed plastic bag for 2 weeks.
- Mattresses should be vacuumed

Alternative Regimes

- Permethrin 5% cream (Lyclear Dermal cream). Apply over whole body and wash off after 12 hours. Repeat at 7 days. Permethrin is not licensed for those under 18years of age. Give 30-60g.

Partner notification

- Advise treatment of current sexual partners
- Partner notification of partners from the previous 3 months

Follow up

- Advise client to re-examine 1 week after final treatment
- Advise client that dead nits may remain adherent to hairs and can be removed with a special comb
- Treatment failures should be given the alternative from the above list, also ensure that application has been correct. Phenothrin is an alternative but the alcohol base may be irritant to the genital area

BNF prices as at Aug 15

Malathion 0.5%	50ml - £4.26 (100ml required)
Permethrin 5%	30g - £5.73 (give 30-60mg)

References

BASHH (2007) United Kingdom Natural Guideline on the management of Phthirus Pubis infestation. Clinical Effectiveness Group. Accessed Feb 19
BNF online accessed Aug 17

<http://www.cks.nice.org.uk/pubic-lice> accessed Feb 19

WOS PUBIC LICE PROTOCOL	APPROVED
WOS MCN CLINICAL GUIDELINES GROUP	VERSION: DRAFT 6.1 LAST UPDATED: Feb 19
REVIEW DATE: Feb 2021 PAGE NUMBER: 2 of 2	COPIES AVAILABLE: www.wossexualhealthmcn.org